

L-S 50 - lincomycin hydrochloride and spectinomycin sulfate tetrahydrate powder, for solution
Pharmacia and Upjohn Company

DIRECTIONS FOR USE

INDICATIONS AND USAGE

For use in chickens up to 7 days of age as an aid in the control of: Airsacculitis caused by either *Mycoplasma synoviae* or *Mycoplasma gallisepticum* susceptible to lincomycin-spectinomycin. Complicated Chronic Respiratory Disease (Air Sac Infection) caused by *Escherichia coli* and *M. gallisepticum* susceptible to lincomycin-spectinomycin.

DOSAGE AND ADMINISTRATION

Dosage

Provide 2 grams (g) antibiotic activity per gallon of drinking water. Administer as the sole source of water for the first 5 to 7 days of life.

Administration

<i>Amount of L-S 50</i>	<i>Amount of Drinking Water</i>
1 bottle	25 gallons
For proportioners delivering 1 ounce of solution per gallon of drinking water, dissolve contents of 5 bottles in each gallon of proportioner solution.	

Important

Chickens should consume water at the following approximate rate to insure intake of the required dose of lincomycin-spectinomycin indicated:

Broilers and Layer Replacements (Light and Heavy)

<i>Age (Weeks)</i>	<i>Daily Water Intake Gallons/1000</i>	<i>Dosage mg Antibiotic/lb</i>
1	5	50-65

PRECAUTION

Discard medicated drinking water daily and replace with fresh medicated drinking water.

WARNING

Not for human use.

Store at Controlled Room Temperature 20° to 25° C (68° to 77° F).

Restricted Drug (California)-Use Only as Directed

Made in China

Distributed by: Pharmacia & Upjohn Company

Div. of Pfizer Inc, New York, NY 10017

PRINCIPAL DISPLAY PANEL - BOTTLE LABEL

L-S 50

WATER SOLUBLE® POWDER

lincomycin-spectinomycin

Antibacterial and Anllmycoplasmal

TAKE TIME

OBSERVE LABEL

DIRECTIONS

FOR USE IN CHICKENS ONLY

This bottle contains:

Lincomycin hydrochloride, equivalent to lincomycin

16.7 grams

Spectinomycin sulfate tetrahydrate, equivalent to spectinomycin

33.3 grams

Total Antibiotic Activity

50.0 grams


NADA #46-109, Approved by FDA

PEEL FROM THE RIGHT


FOR SPANISH LANGUAGE

NET WT. 2.65 OZ


(75.0 grams)



LS 50
WATER SOLUBLE® POWDER
lincomycin-spectinomycin
Antibacterial and Antimycoplasmal



PEEL FROM THE RIGHT
FOR SPANISH LANGUAGE



INDICATIONS AND USAGE: For use in chickens up to 7 days of age as an aid in the control of: Airsacculitis caused by either *Mycoplasma synoviae* or *Mycoplasma gallisepticum* susceptible to lincomycin-spectinomycin. Complicated Chronic Respiratory Disease (Air Sac Infection) caused by *Escherichia coli* and *M. gallisepticum* susceptible to lincomycin-spectinomycin.

DOSAGE AND ADMINISTRATION
Dosage: Provide 2 grams (g) antibiotic activity per gallon of drinking water. Administer as the sole source of water for the first 5 to 7 days of life.

Administration:

Amount of L-S 50	Amount of Drinking Water
1 bottle	25 gallons

For proportioners delivering 1 ounce of solution per gallon of drinking water, dissolve contents of 5 bottles in each gallon of proportioner solution.

Important: Chickens should consume water at the following approximate rate to insure intake of the required dose of lincomycin-spectinomycin indicated:
Broilers and Layer Replacements (Light and Heavy)

Age (Weeks)	Daily Water Intake Gallons/1000	Dosage mg Antibiotic/lb
1	5	50-65


1454000
A73167-1

PRECAUTION: Discard medicated drinking water daily and replace with fresh medicated drinking water.


WARNING: Not for human use. Store at Controlled Room Temperature 20° to 25° C (68° to 77° F).

Restricted Drug (California)—Use Only as Directed

Made in China
Distributed by: Pharmacia & Upjohn Co.
Div. of Pfizer Inc., New York, NY 10017




LOT:
EXP:



LS 50
POLVO HIDROSOLUBLE®
lincomicina-espectinomicina
Antibacteriano y antimicoplásmico



PEEL FROM THE RIGHT
FOR SPANISH LANGUAGE



INDICACIONES Y USO: Para usar en pollos de hasta 7 días de edad, con el fin de ayudar a controlar: La alveolitis causada por *Mycoplasma synoviae* o *Mycoplasma gallisepticum* susceptible a la lincomicina-espectinomicina. Enfermedad Respiratoria Crónica Complicada (Infección de los alvéolos) causada por *Escherichia coli* y *M. gallisepticum* susceptible a la lincomicina-espectinomicina.

DOSIS Y ADMINISTRACIÓN
Dosis: Proporcionar 2 gramos (g) de actividad antibiótica por cada galón de agua de bebida. Administrese como fuente única de agua durante los primeros 5 a 7 días de vida.

Administración:

Cantidad de L-S 50	Cantidad de agua de bebida
1 frasco	25 galones

Para los surtidores que entregan 1 onza de solución por cada galón de agua potable, disuelva el contenido de 5 frascos en cada galón de solución del surtidor fraccionador.

Importante: Los pollos deben consumir agua en las siguientes cantidades aproximadas, para asegurar la ingestión de la dosis requerida de la lincomicina-espectinomicina indicada:
Pollos de carne (broiler) y reemplazo de ponedoras (aves livianas y pesadas)

Edad (semanas)	Ingestión diaria de agua (galones/1000)	Dosis (mg antibiótico/lb)
1	5	50-65

PRECAUCIÓN: Deseche el agua de beber restante todos los días y cámbiela por agua de beber fresca que contiene el medicamento.

ADVERTENCIA: No usar en seres humanos. Almacénese a temperaturas controladas entre 20° y 25° C (68° a 77° F). Fármaco restringido (California)—Úsese solamente según las indicaciones.

Made in China Dist. by: Pharmacia & Upjohn Co., Div. of Pfizer Inc., New York, NY 10017

1454000
A73167-1